



1 **Sectoral Vulnerability to Drought: Exploring the Role of Blue**
2 **and Green Water Dependency in Mid and High-Latitudes**

3

4 Elin Stenfors¹, Malgorzata Blicharska², Thomas Grabs¹, Claudia Teutschbein¹

5

6 1 Uppsala University, Department of Earth Sciences, Program for Air, Water and Landscape Sciences, Villavägen 16, 75236
7 Uppsala, Sweden

8 2 Uppsala University, Department of Earth Sciences, Program for Natural Resources and Sustainable Development,
9 Villavägen 16, 75236 Uppsala, Sweden

10

11 *Correspondence to:* Elin Stenfors (elin.stenfors@geo.uu.se)



12 **Abstract.**

13 The European continent has experienced several large-scale drought events in recent years, and climate
14 projections suggest an increasing drought risk in many parts of the world. As droughts can have large impacts on
15 socio-hydrological systems, analysing drought risk is an important part for proactive drought risk management
16 and disaster risk reduction. Drought risk can be expressed as a product of hazard, exposure and vulnerability,
17 where vulnerability is highly contextual and complex. As droughts can affect all parts of the hydrological system,
18 from precipitation and soil moisture to groundwater and surface water reservoirs, drought vulnerability differs
19 depending on what part of the system is studied. Building on previous results from a survey analysing drought
20 vulnerability across seven water-dependent sectors, this paper explores how vulnerability factors vary based on
21 sectors' dependency on blue water (surface and subsurface freshwater) or green water (soil moisture) in mid and
22 high latitude regions. The findings reveal that drought vulnerability differs based on water type dependency,
23 especially concerning water supply and species characteristics. Perceptions of vulnerability factors vary in
24 number, category, and overall ranking, highlighting the importance of considering water dependency when
25 choosing vulnerability factors for drought risk assessments and to clearly define the drought hazard types
26 involved.



1 Introduction

Climate projections suggest that drought risk is increasing, with some regions likely to experience more frequent and severe drought events in future climates (UNDRR, 2021). Some areas are already seeing more intense drought events, especially West Africa and southern Europe (UNDRR, 2021). The European continent has experienced several large-scale drought events in recent years, with the most recent extraordinary drought conditions in 2022 (Faranda et al., 2023). Droughts can have severe and far-reaching impacts on societies, where impacts can be both direct and indirect (Blauhut et al., 2016), and develop slowly over time, affecting many parts of society (Mishra and Singh, 2010). Among natural hazards, drought events can cause some of the highest economic losses (Kim et al., 2015), while their (often substantial) impacts on human health and the environment are more difficult to assess (UNDRR, 2019).

Modern approaches to drought risk management follow the Sendai Framework for Disaster Risk Reduction (2015–2030). The Sendai framework emphasizes the need for disaster risk management, and aims at furthering the understanding of complex disaster risk. The framework divides risk into three dimensions – hazard, exposure and vulnerability (UNDRR, 2015). A disaster occurs when a hazardous event takes place in a vulnerable area with exposed entities creating a serious disruption of the functions of a community or society (UNDRR, 2019). Hence, in order to understand and manage drought risk, an understanding of all three dimensions is needed.

As a *hazard*, droughts are referred to as creeping phenomena, with a slow onset and where impacts can appear after the drought event itself has ended. As a result, determining its onset and termination can be difficult. Droughts are generally distinguished as four different types – meteorological droughts that represent a lack of precipitation, sometimes combined with an increase in evapotranspiration compared to normal conditions (Van Loon, 2015), agricultural drought depicting a soil moisture deficit that affects soil vegetation and crops (Van Loon, 2015), hydrological drought that is characterized by deficits in surface- and/or subsurface water resources (Mishra and Singh, 2010), and socioeconomic drought that represents an impact-oriented deficit of water as a good in the supply and demand network of the human-water nexus (Mishra and Singh, 2010). The different drought types have different propagation times, where meteorological and soil moisture drought typically develop faster than hydrological or socioeconomic droughts, as reservoirs can off-set and smooth out effects of a drought over time (Mishra and Singh, 2010; Van Loon, 2015).

The second risk dimension, drought *exposure*, encompasses the entities exposed during a drought event, such as buildings, inhabitants, or crops (Ciurean et al., 2013). The exposure component is often expressed as a sub-dimension of vulnerability, as the term was originally included in the definition of vulnerability in the older IPCC Assessment Report (AR) 4 (IPCC, 2007). However, the exposure term has since then been omitted from the vulnerability definitions used in AR5 and AR6 (IPCC, 2022; IPCC et al., 2014) and is now considered a standalone dimension of drought risk. Instead, the third risk dimension, drought *vulnerability*, can generally be expressed as the predisposition of a system to be negatively affected by a drought (Füssel, 2007). Yet, vulnerability is a complex concept, which can be described, defined and conceptualized in a number of ways (Adger, 2006; Ciurean et al., 2013; Sebesvari et al., 2016; Turner et al., 2003). In IPCC AR6 vulnerability is defined as encompassing “a variety of concepts and elements including sensitivity or susceptibility to harm and lack of capacity to cope and adapt” (IPCC, 2022). Adaptive capacity can be seen as the ability to anticipate and learn from droughts in the long term, whereas coping capacity refers to the ability to react and cope in the short term. Susceptibility on the other hand relates to the predisposition to be negatively impacted by a drought event. Using the definition proposed by IPCC AR 6 (IPCC, 2022), Stenfors et al. (2024a) conceptualized drought vulnerability of a socio-



hydrological system as comprising three dimensions: (1) governance processes and the available policies and plans (i.e., governance), (2) the indirect water consumers of the system (i.e., society), and (3) its direct water consumers (i.e., particular sectors) (Figure 1a). Factors of vulnerability connected to governance processes, policies and plans, can affect both drought vulnerability in individual sectors, as well as society as a whole. They relate to aspects such as drought awareness of authorities, presence of drought plans and risk assessments, or the financial capacity of the government to offer support during a drought event. Meanwhile, vulnerability factors connected to indirect water consumers relate to the population consuming water indirectly, through goods such as food, energy or public water supply. For indirect water consumers, vulnerability can be expressed through demographic aspects that may affect drought vulnerability, such as different aspects of socio-economic susceptibility (e.g., level of income or integration). Lastly, direct water consumers refer to socio-hydrological sectors that consume water directly for production of goods or for sustaining ecosystem services.




Figure 1. (a) The conceptual framework developed by Stenfors et al (2024a). In the framework drought vulnerability is represented by factors relating to governance, indirect water consumers and direct water consumers. The latter is further divided into factors relating to sectors dependent on blue or green water respectively. The framework has been modified from Stenfors et al. (2024) to visualize how different drought types (i.e., hydrological and soil moisture droughts) are linked to different direct water consumers. (b) Geographic extent of survey participants representing water-dependent sectors in Sweden.

These direct water consumers can be further categorized based on their dependency on different types of water resources, i.e., whether they depend predominantly on *green* or *blue* sources of water. Falkenmark and Rockström (2006) described green water as water stored in the unsaturated zone as soil moisture, whereas blue water is stored as sub-surface water in the saturated zone or as surface water. Examples of direct water consumers relying on surface or subsurface water (i.e., *blue water resources*) include energy producers, drinking water suppliers, irrigated agriculture and aquatic ecosystems. In contrast, sectors like forestry, rainfed agriculture and terrestrial ecosystems, primarily depend on water stored in the unsaturated zone as soil moisture (i.e. *green water*).



93 The distinction between green and blue water resources is crucial as they are affected differently by various types
 94 of drought. Agricultural drought represents deficits in green water supplies, whereas hydrological drought refers
 95 to water deficits in blue water. Hence, the conceptual framework of Stenfors et al (2024a) suggests that different
 96 drought types could affect various consumers/sectors differently, each with their own specific vulnerabilities to
 97 that drought type. To explore these consumer/sector-specific vulnerabilities, Stenfors et al. (2024b) conducted an
 98 online survey targeting stakeholders from seven water-dependent sectors in Sweden, northern Europe (Figure 1b).
 99 Stretching from 55°2'N to 69°3'N, Sweden comprises three climate zones according to the Köppen-Geiger
 100 classification, namely warm-summer hemi boreal (Dfb) in the south, subarctic boreal (Dfc) in central and northern
 101 Sweden, and tundra (ET) in the north-west, where many areas currently classified as Dfb and Dfc are projected to
 102 transition to Cfb and Dfb by the end of the century (Beck et al., 2018). Sweden has historically been considered
 103 water-abundant, however the 2018 drought affected large parts of the country (Teutschbein et al., 2022).
 104 The survey results enabled the identification of user-validated drought vulnerability factors that are particularly
 105 relevant to water-dependent sectors and societies in mid and high-latitude regions (located above 50°N). The
 106 diverse perceptions of drought vulnerability found across different consumers/sectors further indicate that
 107 variations in the relevance of vulnerability factors may stem from the specific type of water dependency (green
 108 versus blue) of each consumer/sector.

109 Building on the notion of an inherent relation between sectoral drought vulnerability and the type of water
 110 dependency in mid and high latitude regions (Stenfors et al., 2024b), this paper provides a comprehensive analysis
 111 of how sectoral drought vulnerability differs according to water-type dependency, and discusses the implication
 112 for drought risk assessments, as well as policy design. Utilizing the conceptual framework described by Stenfors
 113 et al (2024a) and the survey data presented by Stenfors et al. (2024b) in this issue, we aim to address the following
 114 research questions for drought vulnerability in mid and high latitude regions:

- 115 1. What are the relevant vulnerability factors for blue water-dependent sectors compared to green water-
 116 dependent sectors?
- 117 2. How do the vulnerability factors for blue and green water-dependent sectors rank in terms of impact
 118 scores?
- 119 3. Which vulnerability factors are rated the highest for each type of water dependency?
- 120 4. Do the impact ratings of vulnerability factors vary among respondents based on the type of water
 121 dependency?

122 **2 Methods and Data**

123 **2.1 Data collection**

124 The starting point for the analysis was a list of 74 vulnerability factors connected to direct water consumers
 125 compiled from existing literature by Stenfors et al (2024a) for forested cold climate regions. These factors related
 126 broadly to nine categories of vulnerability (Figure 2) and referred to adaptive capacity, coping capacity or
 127 susceptibility - following the IPCC AR6 definition of vulnerability.



No. of factors	Categories of vulnerability	Dimensions
4	Irrigation: Presence of irrigation	C
5	Funds: Available funds and financial capacity	C S
6	Authority: Characteristics of authority	C
8	Stress: Presence of anthropogenic stress	S
9	Policies: Presence of policies and plans	A C
9	Setting: Conditions of surrounding setting	S
9	Species: Species characteristics	A S
10	Supply: Water supply	A C S
14	Tools: Availability of tools and resources	A C

Figure 2. Overview of the number of vulnerability factors for direct water consumers compiled by (Stenfors et al. 2024). The factors are divided into nine subcategories (short names in bold) relating to the general attributes of the factors. Information as to whether each category includes factors connected to adaptive capacity (A), coping capacity (C) and/or susceptibility (S) is also displayed.

To validate and refine the assembled vulnerability factors, an online survey was developed, targeting seven water-dependent sectors in Sweden (Figure 1b) – energy production including for example hydropower or nuclear production (i.e., energy), agricultural crop production or livestock (agriculture), aquatic and terrestrial ecosystems (environment), drinking water production and distribution (water supply), water resource management (water resources), forest conservation and production (forestry), and water intensive industries such as paper and pulp, chemical production, or steel and metal works (water intensive industry). The survey included three sections:

- I. Background Information:** This section collected data about the respondents, including their primary sector, years of experience, type of organization and Swedish region in which they work.
- II. Sector-Specific Vulnerability Factors:** Respondents were asked to rate the importance of various vulnerability factors for drought risk in their primary sector, using a 5-point scale (0 = no impact to 4 = high impact)
- III. Societal Vulnerability Factors:** Respondents rated the importance of vulnerability factors for drought risk across Swedish society, again using the 5-point rating scale.

For a more detailed description of the survey, we refer the reader to Stenfors et al. (2024b) and the supplementary information (S1, section S1.1)

The survey was sent to 561 recipients in spring 2023, including governmental/local authorities (354 recipients), private/municipal/state-owned enterprises (81), academia or research institutes (46), national/regional trade associations (45), and NGOs (35). Receiving 108 responses in total, the response rate was 19.3%. Initial data cleaning identified six respondents answering exclusively “I don’t know”, and their responses were excluded from all subsequent analysis. Furthermore, one respondent did not specify a sectoral focus and could therefore not be included in the analysis. A more detailed description of the recipient selection process can be found in the supplementary materials (S1, section S1.2).

2.2 Data Analysis

Survey responses were analyzed in Microsoft Excel and RStudio using a four-step approach involving data preparation, relevant factor identification, vulnerability factor ranking and hypothesis testing.



158 As a first step, survey responses were categorized based on the primary sector into three groups: blue water-
 159 dependent, green water-dependent, or universal water consumers (i.e. relying on both blue and green water).
 160 Respondents from the energy, drinking water and water resource sector were categorized as primarily blue water-
 161 dependent. Those in the forestry sector were categorized as mainly green water-dependent. Within the
 162 environmental sector, respondents were further divided into three groups: those working with aquatic ecosystems
 163 were classified as blue water-dependent, while those focused on terrestrial ecosystems were considered green
 164 water-dependent. The third group, referred to as universal water consumers, included respondents working with
 165 both aquatic and terrestrial ecosystems, and respondents from the agricultural sector, as their reliance on either
 166 blue or green water could not be distinctly categorized.

167 Second, relevant vulnerability factors for the two groups respectively were identified following Meza et al. (2019)
 168 and Stenfors et al. (2024b). A vulnerability factor was considered relevant for the water dependent groups if 50%
 169 or more of survey respondents rated it as having a medium high or high impact (i.e., a median score of three or
 170 higher) on drought risk in their sector. Hence, drought vulnerability factors relevant for blue, green and universal
 171 water consumers were identified by calculating the median rating, grouped by water type consumption (blue,
 172 green or universal water consumption).

173 After identifying relevant factors, factor impact scores were calculated in a third step. This allowed for analysis
 174 of the relative importance of vulnerability factors depending on water type dependency. The respondent ratings,
 175 originally ranging from 0 to 4, were rescaled into a scale of 0-1, with 0.25-step increments. The impact score was
 176 then attained by calculating the mean rescaled rating for each factor, grouped by water consumption type. Highly
 177 impactful vulnerability factors have impact scores closer to 1, whereas less impactful factors have impact scores
 178 close to 0. The calculated impact score is not an indication on whether the factor has a positive or negative impact
 179 on drought risk, only that the factor is perceived as impactful.

180 Lastly, pairwise-Wilcoxon rank sum tests with corrections for multiple testing were used for analyzing significant
 181 differences in ratings between the three water consumer groups, using pairwise comparisons between the three
 182 water consumptions groups (i.e. blue, green and universal water consumption).

183 **3 Synthesis of Results**

184 **3.1 Respondent characteristics**

185 Grouping the respondents based on sectoral water type dependency resulted in three groups, where blue water
 186 consumers had the largest number of respondents (48 respondents), followed by universal water consumers (29),
 187 and green water consumers (24 respondents) (Table 1). In all three groups, a majority of the respondents were
 188 working in authorities (60% across all three groups), followed by respondents working in research (19% in total)
 189 and private/municipal/authority owned enterprises (12%). The blue water consumers, as well as universal
 190 consumers both had respondents working for an NGO and trade associations, where the universal water consumers
 191 had five respondents working for agricultural trade associations. Most respondents were located in southern
 192 Sweden (84%) and indicated to have a significant experience of working with drought in their primary sector
 193 (56%). More information on respondents' characteristics is presented in the supplementary materials (S2).

194




195 **Table 1. Overview of respondent characteristics, divided by water consumption type (blue, green, universal), place of**
 196 **employment, geographical location in Sweden, their reported experience with drought, as well as their primary sectoral**
 197 **focus.**

<i>Organization</i>	<i>Blue</i>	<i>Universal</i>	<i>Green</i>	<i>Grand total</i>
<i>Authority</i>	31	16	14	61
<i>Enterprise</i>	7	1	4	12
<i>NGO</i>	1	1		2
<i>Research</i>	7	6	6	19
<i>Trade association</i>	2	5		7
<i>Location</i>				
<i>North</i>	6	6	4	16
<i>South</i>	42	23	20	85
<i>Drought experience</i>				
<i>Limited</i>	8	7	1	16
<i>Moderate</i>	13	7	8	28
<i>Significant</i>	27	15	15	57
<i>Sector</i>				
<i>Agricultural</i>		13		13
<i>Energy</i>	7			7
<i>Environmental</i>	9	16	10	35
<i>Forestry</i>			14	14
<i>Water intensive industry</i>	1			1
<i>Water resources</i>	15			15
<i>Water supply</i>	16			16
<i>Respondents (number)</i>	48	29	24	101

198 3.2 Relevance of vulnerability factors for blue, green and universal water consumers

199 In total 63 vulnerability factors were considered relevant by at least one of the three water consuming groups
 200 (Figure 3a). Universal water consumers, reliant on both blue and green water, found the largest number of factors
 201 as relevant (60 factors), followed by blue water consumers (43), and green water consumers (18). Universal water
 202 consumers found vulnerability factors relating to all nine categories relevant for their sectors. This can be
 203 compared to green water consumers, who only found factors relating to five categories as relevant. The factors
 204 relevant for green water consumers were mainly related to species characteristics, conditions of the surrounding
 205 settings and available tools & resources. The respondents in this group also found one factor related to policies
 206 and one related to anthropogenic stress as relevant. On the other hand, blue water consumers found at least one
 207 factor as relevant in all factor categories except one. No factors related to species characteristics were considered
 208 relevant for blue water consumers. In contrast, blue water consumers found all factors related to policies and plans
 209 relevant, as well a majority of the factors relating to the conditions of surrounding setting, available water supply,
 210 and available tools and resources.



211

212 **Figure 3. (a) Number of vulnerability factors included in the survey for direct water consumers (Survey), the number**
213 **of factors relevant by at least one water consumption group (Relevant), as well as the number factors identified as**
214 **relevant for each water consumptions group. Each factor is categorized as belonging to one out of nine categories. (b)**
215 **ratio and number of factors belonging to adaptive capacity (dark blue), coping capacity (blue), or susceptibility (light**
216 **blue), out of the total number of factors relevant for all respondents, as well as for blue, green and universal water**
217 **consumers respectively.**

218 Universal water consumers found factors related to all three dimensions of vulnerability as relevant (Figure 3b),
219 where the ratio of factors belonging to adaptive capacity, coping capacity or susceptibility was approximately the
220 same as what was included in the survey. Blue water consumers, whilst also finding factors connected to all three
221 dimensions, found a large share of factors relating to coping capacity as relevant for the group. This can be
222 compared to green water consumers, who only found one factor relating to coping capacity as relevant, as most
223 relevant factors for green water consumers related to either susceptibility or adaptive capacity.

224 3.3 Impact scores for vulnerability factors for blue, green and universal water consumers

225 The highest impact scores for several factors were given by universal water consumers (Figure 4). This group
226 gave high impact scores for factors relating to adaptive capacity, coping capacity and susceptibility, however the
227 highest impact scores were given to factors connected to susceptibility. Blue and green water consumers also gave
228 factors relating to susceptibility the highest impact scores. Green water consumers, primarily finding factors
229 relating to susceptibility and adaptive capacity relevant, seemed to find factors relating to susceptibility slightly
230 more impactful than factors concerning adaptive capacity, especially the ones relating to species characteristics
231 or the conditions of the surrounding settings. Conditions of the surrounding settings, policies and available tools
232 and resources, were the three categories of factors that included factors found relevant for all three water
233 consuming groups. Examples of factors found relevant for all three consumer groups were the soil water holding
234 capacity, presence of wetlands, lakes, and ponds, and the availability of drought risk assessments. In general, blue
235 water consumers gave high impact scores to factors relating to water supply, policies and plans, or conditions of
236 the surrounding settings. Similarly, universal water consumers gave many of their high impact scores to factors
237 relating to water supply and the conditions of the surrounding settings, followed by anthropogenic stress. Whilst
238 also giving high impact scores to factors concerning the conditions of the surrounding settings, green water



consumers also gave factors relating to species characteristics and available tools and resources some of their highest impact scores. Relevant vulnerability factors and their corresponding impact scores for the three consumers groups are available in the supplementary materials (S3).





Figure 4. Impact scores for vulnerability factors concerning adaptive capacity (adaptive), coping capacity (coping), and susceptibility, rated regarding their impact on drought risk in blue, green and universal water-dependent sectors. Filled dots indicate that the factor is considered relevant for the consumers (i.e., with a median score of 3 or higher), whereas open circles indicate that the factor is not considered relevant. The point size signifies the percentage of respondents within a consumer group that provided an impact rating for the factor.



248 **3.4 Highest rated vulnerability factors for blue, green and universal water consumers**
249 The top highest rated vulnerability factors for green water consumers, related to species characteristics or the
250 conditions of the surrounding settings (Figure 5). Their highest rated factors were soil water holding capacity,
251 followed by the drought tolerance of current species. For blue water consumers, two of the three highest rated
252 factors related to conditions of the surrounding settings, such as the presence of wetlands, lakes and ponds and
253 the geographical characteristics of the area. The second highest rated factor for blue water consumers was baseline
254 water stress. When comparing the highest rated vulnerability factors for universal water consumers with the other
255 two groups, it can be seen that universal water consumers shared several factors with either green or blue water
256 consumers. For example, their top-rated factor, baseline water stress, was the second highest rated factor for blue
257 water consumers. The second highest vulnerability factor for universal consumers, the soil water holding capacity,
258 was the highest rated factor among green water consumers.



259
260 **Figure 5. The highest rated factors divided per water consumer group.**



3.5 Differences in impact scores depending on blue, green and universal water dependency

The three groups of water consumers showed different patterns in their ratings of the nine categories of factors (Figure 6). For example, blue water consumers, showed a relatively large spread in their ratings for factors concerning, for example, conditions of the surrounding settings and anthropogenic stress. On the other hand, blue water consumers consistently rated factors concerning species characteristics low, and policies and plans high. Contrarily, universal consumers tended to rate most factors highly, where the largest spread in ratings was seen for factors relating to water supply and anthropogenic stress. The universal water consuming group included agriculture, which was reflected in the results, as factors relating to irrigation received higher impact scores by the group compared to blue and green water consumers. Green water consumers generally rated most categories of factors lower than the other consumer groups, where the largest spread in impact scores was for factors concerning conditions of the surrounding settings. The group consistently gave low impact scores for factors concerning water supply, irrigation and available funds and financial capacity.





Figure 6. Distributions of impact scores for the nine categories of vulnerability factors for blue water consumers (left), green water consumers (right) and universal water consumers (middle).

Universal water consumers shared several factors with high impacts scores with both blue and green water consumers respectively (Figure 7). Both universal and blue water consumers, highly rated several factors connected to water supply, policies and plans, anthropogenic stress, available tools and resources, and characteristics of authorities. In comparison, universal and green water consumers, shared fewer highly rated factors. Universal and green water consumers mainly shared high impact scores for factors relating to species characteristics, conditions of the surrounding settings, and available tools and resources. Furthermore, the two groups rated a few factors relating to the characteristics of authority, and policies and plans highly.



283
284
285 **Figure 7. Impact scores for 63 drought vulnerability factors rated by universal water consumers (y-axis), blue water**
286 **consumers (x-axis, left) and green water consumers (x-axis, right). Colors indicate that the factors belong to one out of**
287 **the nine factor categories. Observe that the x- and y-axis has been adjusted for better data visualization, and does not**
288 **start at zero. The thicker grey lines, mark the threshold above which vulnerability factors have a medium high to high**
289 **impact score. This plot, with accompanying text labels for the individual factors, can be found in supplementary**
290 **materials (S4 section S4.1 and S4.2).**

291 In general, blue water consumers rated a majority of factors slightly higher than green water consumers (Figure
292 8). However, green water consumers rated factors concerning species characteristics highly, as well as a majority
293 of factors relating to the conditions of surrounding settings. In contrast, all factors concerning water supply
294 received low impact ratings by the group, as well as factors relating to irrigation and available funds. Conversely,
295 blue water consumers rated factors relating to water supply and policies highly, whilst all factors relating to species
296 characteristics received low impact scores. Several factors relating to available tools and the conditions of the
297 surrounding settings received high impact scores from both blue and green water dependent sectors. Looking at
298 the highest rated factors, common for the two groups, factors were mainly related to available tools, the conditions
299 of the surrounding settings and policies and plans.




Figure 8. Impact scores for 63 drought vulnerability factors rated by blue water consumers (x-axis) and green water consumers (y-axis). Colors indicate that the factors belong to one out of the nine factor categories. Observe that the x- and y-axis has been adjusted for better data visualization, and does not start at zero. The thicker grey lines, mark the threshold above which vulnerability factors have a medium high to high impact score. This plot, with accompanying text labels for the individual factors, can be found in supplementary materials (S4, section S4.3).

When comparing ratings between green and blue water consumers, significant differences were seen for factor ratings for one or more factors in seven out of nine factor categories (Table 2). No significant differences between ratings made by blue and green water consumers were seen for factors relating to authority and tools. However, significant differences were seen for nine out ten factors relating to water supply. When comparing factor ratings for universal and blue water consumer groups, significant differences were seen for several factors relating to species characteristics, whereas no significant differences were seen for factor ratings relating to authorities, policies and plans, anthropogenic stress, and available tools and resources. Three factor categories stood out as having significant differences when comparing factor ratings made by universal water consumers and green consumers, namely funds, irrigation and water supply. For example, significant differences were seen for factor ratings for all four factors relating to irrigation, and seven out of ten factors relating to water supply, for these two groups. All factors exhibiting significant differences in rating between the consumers groups and their corresponding p-values are presented in the supplementary materials (S5).



Table 2. Number of factors per factor category with significant differences in ratings between respondents from the three water consumption groups, together with the number of factors included in each of the nine categories. Significant differences identified using pair-wise Wilcoxon rank sum tests are shown for the water consumption pairs; green and blue water consumers (Green-Blue), blue and universal water consumers (Universal-Blue), and green and universal water consumers (Universal-Green).

	Survey	Green-Blue	Universal-Blue	Universal-Green
Authority	6	0	0	0
Funds	5	2	1	3
Irrigation	4	4	1	4
Policies	9	2	0	0
Setting	9	1	2	2
Species	9	4	5	0
Stress	8	3	0	3
Supply	10	9	1	7
Tools	14	0	0	0
Grand Total	74	25	10	19

329

330 4 Discussion

331 Our study explored the role of water dependency for drought vulnerability within a socio-hydrological system.
 332 Proactive drought risk management requires an integrated analysis of drought hazard, exposure and vulnerability.
 333 Using user validation to analyse drought vulnerability across sectors categorized by their dependency on blue
 334 and/or green water sources provided a unique opportunity to deepen our understanding of drought vulnerability
 335 with respect to drought type and water dependency. This approach also enhances the quality of future drought risk
 336 assessments. We found notable differences in the perceived relevance, impact scores and overall ratings of various
 337 drought vulnerability factors among blue, green and universal water consumers.

338 Universal water consumers found the largest number of relevant factors. This outcome aligns with the distinction
 339 between blue and green water dependency, as universal consumers are likely to consider factors related to both
 340 types of water sources as relevant. In fact, all factors that were found relevant by either blue or green water
 341 consumers, were also relevant for universal consumers, with the exception for three vulnerability factors that were
 342 only considered relevant by blue water consumers. Notably, blue water consumers identified more than twice as
 343 many relevant factors as green water consumers, many of which concerning water supply or the availability of
 344 tools and resources for managing drought and water availability. This could potentially be due to many factors
 345 found in the literature review being more geared towards vulnerability of blue water consumers. In fact , in the
 346 survey, 17 factors were directly related to blue water, its governance, related policies, or monitoring tools, while
 347 only 12 factors were related to species characteristics, forest management practices or vegetation modelling.
 348 Although this unequal focus in the survey might partially contribute to the difference in the total number of
 349 relevant factors between the two groups, it cannot fully explain the wide gap and indicates that there are indeed
 350 underlying differences in how drought vulnerability is perceived between blue and green water consumers.

351 Our results highlight that blue and universal water consumers found several factors related to policies and plans,
 352 such as having authority level drought management plans, local water management plans and planned drought
 353 prevention measures as relevant for drought risk. Furthermore, the two groups found the existence of water use
 354 priority classes and defined water use rights as relevant. This validates the importance of incorporating drought
 355 and water management plans as tools for increasing adaptive and coping capacity for blue water consumers in



socio-hydrological systems, as has been described and promoted in literature and international declarations (Sivakumar et al., 2014; UNDRR, 2021; Wilhite et al., 2014). However, further research is needed to better understand the role of drought management plans for green water consumers, as only one vulnerability factor connected to policies and plans was deemed relevant for the group, namely the presence of a local water management plan. Notably, less than half of the respondents in this group provided a rating for this factor. Other policy-related factors, such as the existence of an authority-level drought management plan or planned drought prevention measures at the authority level, were not considered relevant by green water consumers. The reasons for this lack of relevance are unclear and warrant further investigation to better understand how drought-related policies can be adapted to be more applicable to green water consumers. Nevertheless, this may imply that there is a gap between policy tools and this consumer group, that need to be better analyzed for improved policy support for green water consumers.

Instead, green water consumers found seven out of nine factors relating to species characteristics to be relevant. This category included factors such as drought tolerance of species and root depths, which considerably influence the effects of green water deficits. Only green and universal water consumers considered this category relevant, likely due to the minimal impact these factors have on blue water consumers such as energy production and water supply during a hydrological drought, apart from the competing water needs arising from potentially irrigating the species. Factors such as competing water needs, and other factors concerning anthropogenic stress on water resources were instead considered relevant for blue and universal water consumers.

Both species characteristics and anthropogenic stress are categories of factors relating to susceptibility. Whilst the three water consumer groups considered different factors concerning susceptibility relevant, this dimension of vulnerability was generally relevant for all three groups. The same can be said for adaptive capacity, where all three consumer groups found several relevant factors. Factors relating to coping capacity saw varying relevance among the consumer groups, where the largest number of relevant factors for blue water consumers belong to this dimension. In contrast, green water consumers found only one factor relating to coping capacity to be relevant, which might indicate that this group has limited tools for coping with drought events. As a result, they may place greater emphasis on anticipatory approaches that focus on increasing adaptive capacity and decreasing susceptibility. However, this could also be a result of the specific vulnerability factors that were categorized as coping capacity in the survey, as many focused on financial capacity, policies and characteristics of governance, rather the reactionary management measures that can be employed in the case of droughts. For example, measures for mitigating drought effects on forests, such as thinning, can be both anticipatory and reactionary. In the survey, such factors were not included as standalone factors, but were incorporated in the factor “Use of adaptive measures”. Similarly, specific management measures for forestry and agriculture were subject to the same aggregation.

In total, ten vulnerability factors were considered relevant for all three water consumer groups. The factors primarily related to two categories of vulnerability, i.e., the conditions of the surrounding settings and the available tools and resources. The list includes factors such as the availability of a drought risk assessment and having access to relevant data regarding drought, as well as aspects such as the soil water holding capacity and the



396 presence of wetlands, lakes, and ponds. Hence, these factors could potentially be viewed as universal vulnerability
 397 factors, relevant regardless of water type dependency or drought type exposure, suggesting that certain baseline
 398 conditions and resources are critical for managing drought risks, irrespective of the specific type of water
 399 dependency or drought exposure. This universality implies that these factors are foundational to overall resilience
 400 against drought, serving as key elements that all sectors must address to reduce vulnerability. This further supports
 401 the argument for cross-sectoral collaboration in drought preparedness and response strategies (e.g., Bretan and
 402 Engle, 2017; Medel et al., 2020), ensuring that all sectors can benefit from shared tools and data.

403

404 Both similarities and differences were found when studying the distribution of impact scores. All three consumers
 405 groups generally gave factors relating to the conditions of the surrounding settings high impact scores.
 406 Furthermore, each water consumer group had at least one factor concerning the conditions of the surrounding
 407 setting among its five factors with highest impacts scores. The attributed impacts scores for this factor category,
 408 however, varied greatly among the respondents for several factors. Whilst blue and universal consumers both gave
 409 high scores to several factors related to water supply, blue water consumers put greater emphasis on policies and
 410 plans compared to universal consumers.

411

412 Universal water consumers gave some of the highest impacts scores among the three consumers groups. Baseline
 413 water stress, the soil water holding capacity and having a reliable water resource for water supply received the
 414 highest impacts scores from universal water consumers. Interestingly, these factors also received high impact
 415 scores from blue water consumers, with water stress and reliable water resources ranking among the highest for
 416 them as well. Soil water holding capacity received high impact scores from all three consumers groups, but it
 417 stood out as one of the top factors for both universal and green water consumers. Across all three water consumer
 418 groups, the factors with the highest impact scores primarily concerned susceptibility, suggesting that minimizing
 419 susceptibility is a main priority for all water consumer groups.

420

421 It is worth noting that some factors connected to irrigation, such as the amount of water available for irrigation
 422 and the use of effective irrigation systems, received higher impact scores from blue water consumers compared
 423 to green water consumers. While the exact reason for this is unknown, we can speculate that irrigation can put
 424 additional pressure on water resources, leading to competing water interests among blue water consumers. Such
 425 competition can exacerbate water stress during drought events in areas where total consumption needs exceed water
 426 supply (Famiglietti, 2014; Rossi, L., et al., 2023). For example, in their drought vulnerability analysis for Finland,
 427 Ahopelto et al (2019) found that some areas in southern Finland would have difficulties in supplying water for
 428 the calculated consumptive water needs during a simulated severe drought. Universal water consumers also found
 429 factors concerning irrigation as impactful. This may partly result from the inclusion of respondents from the
 430 agricultural sector, as Stenfors et al (2024b) noted that this sector rated such factors particularly highly.

431 Significant differences in the ratings of vulnerability factors based on water dependency were apparent for factors
 432 relating to species characteristics, irrigation and water availability and supply. This implies that different water-
 433 consuming sectors exhibit different vulnerabilities. Green water consumers tend to focus primarily on
 434 vulnerability factors that affect or are affected by soil moisture deficits, such as drought-tolerant species and the
 435 soil water holding capacity. Meanwhile, blue water consumers are more concerned with aspects related to the



availability, regulation and use of blue water resources, including the reliability of water resources for water supply, authority level water strategies, and the presence of water stress. Consequently, as anticipated, blue water consumers mainly focus on vulnerability to blue water deficits and green water consumers on vulnerability to green water deficits. These distinctions have implications for drought risk assessments, underscoring the need to consider the specific type of drought hazard when designing drought vulnerability and risk assessments. Notably, Hagenlocher et al. (2019) found that 60% of the risk assessments included in their literature review did not specify the drought hazard type used in the assessment. Based on our findings, failing to define the drought hazard type could compromise the quality of the risk assessment, as the use of unsuited vulnerability factors may cause an under- or overestimation of drought risk, depending on the exposed entities involved. Building on the results found by Stenfors et al (2024b), future vulnerability and risk assessments should be designed with caution to ensure that the selected vulnerability factors accurately reflect both the sectors included in the analysis and their specific water dependencies. This consideration will be particularly crucial for holistic approaches that incorporate multiple socio-hydrological sectors in their analysis, each of which may be vulnerable to different drought types, as well as anthropogenic pressures.

However, it is important to acknowledge certain limitations in our study. For instance, respondents for the agricultural sector could not be distinguished between those engaged in rainfed versus irrigated agriculture, which limits insight into potential differences in agricultural drought vulnerability based on water dependency. Furthermore, with only one response from water-intensive industries, their input on drought vulnerability among blue water consumers was minimal. Despite this, the blue-water-consuming group included respondents from various sectors - working with energy, water resources management, aquatic ecosystems and drinking water supply – providing a broad perspective on drought vulnerability. Finally, our analysis was based on the perceived impact of vulnerability factors on drought risk within respondent's individual sectors. The survey did not collect data on whether impacts are perceived as positive or negative, a gap that future research should address to validate and refine the factors for future vulnerability assessments.

5 Conclusion

Using survey data regarding drought vulnerability in seven water dependent sectors, differences in drought vulnerability in relation to water dependency could be explored. The results showed that drought vulnerability differs depending on water type dependency, especially for vulnerability relating to water supply and species characteristics. Differences in the perception of vulnerability factors between the groups were seen both regarding the number of relevant vulnerability factors, as well as the category of factors found relevant. Furthermore, differences in the impact scores given to vulnerability factors depending on water type dependency were seen. The results reaffirms the division suggested by Stenfors et al. (2024a), where drought vulnerability of direct water consumers depends on the water type dependency of the exposed consumer. The results also highlighted factors that seem to be generally impactful for all consumer groups, offering insights into potential universal vulnerability factors, relevant for all water type dependency. The impact of policies and plans on drought vulnerability of blue and universal water consumers, confirms their importance for tackling drought risk in socio-hydrological systems. However, further research is needed to better understand their impact on green water consumers. Drought risk is expected to increase in many areas and drought risk assessments are important tools for producing effective drought risk management strategies to minimize their impacts. Consequently, future drought



476 vulnerability and risk assessments should put emphasis on clearly establishing the frame for the analysis, focusing
477 on careful selection and consideration of vulnerability factors based on the studied drought type, the exposed
478 entities and their specific drought vulnerabilities to that drought type.

479 **6 Data availability**

480 The data used in this study was collected from respondents under a confidentiality agreement. The confidentially
481 agreement assured respondents that data would be anonymized and only be used within the frame of the PhD
482 project conducted by the corresponding author. Hence, with respect to the confidentiality agreement, data cannot
483 be publicly shared. However, an overview of the survey design and recipient selection procedures are available
484 as supplementary materials. Please contact the corresponding author, for any questions regarding the data.

485 **7 Author Contribution**

486 Elin Stenfors was the lead in conceptualization, data collection and curation, formal analysis, validation,
487 visualization, and writing. She was supported by Malgorzata Blicharska and Thomas Grabs in visualization,
488 methodology development and writing. Claudia Teutschbein was the lead in funding acquisition, and supervision,
489 and supported the corresponding author in all project parts, from conceptualization to writing.

490 **8 Competing interests**

491 The authors declare that they have no conflict of interest.



9 References

- Adger, W. N.: Vulnerability, *Global Environmental Change*, 16, 268–281, <https://doi.org/10.1016/j.gloenvcha.2006.02.006>, 2006.
- Ahopelto, L., Veijalainen, N., Guillaume, J. H. A., Keskinen, M., Marttunen, M., and Varis, O.: Can there be water scarcity with abundance of water? Analyzing water stress during a severe drought in Finland, *SUSTAINABILITY*, 11, <https://doi.org/10.3390/su10021548>, 2019.
- Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., and Wood, E. F.: Present and future Köppen-Geiger climate classification maps at 1-km resolution, *Sci Data*, 5, 180214, <https://doi.org/10.1038/sdata.2018.214>, 2018.
- Blauhut, V., Stahl, K., Stagge, J. H., Tallaksen, L. M., De Stefano, L., and Vogt, J.: Estimating drought risk across Europe from reported drought impacts, drought indices, and vulnerability factors, *Hydrol. Earth Syst. Sci.*, 20, 2779–2800, <https://doi.org/10.5194/hess-20-2779-2016>, 2016.
- Bretan, E. and Engle, N. L.: Drought Preparedness Policies and Climate Change Adaptation and Resilience Measures in Brazil: An Institutional Change Assessment, in: *Evaluating Climate Change Action for Sustainable Development*, edited by: Uitto, J. I., Puri, J., and Van Den Berg, R. D., Springer International Publishing, Cham, 305–326, https://doi.org/10.1007/978-3-319-43702-6_17, 2017.
- Ciurean, R., Schroter, D., and Glade, T.: Conceptual Frameworks of Vulnerability Assessments for Natural Disasters Reduction, in: *Approaches to Disaster Management - Examining the Implications of Hazards, Emergencies and Disasters*, edited by: Tiefenbacher, J., InTech, <https://doi.org/10.5772/55538>, 2013.
- Falkenmark, M. and Rockström, J.: The New Blue and Green Water Paradigm: Breaking New Ground for Water Resources Planning and Management, *J. Water Resour. Plann. Manage.*, 132, 129–132, [https://doi.org/10.1061/\(ASCE\)0733-9496\(2006\)132:3\(129\)](https://doi.org/10.1061/(ASCE)0733-9496(2006)132:3(129)), 2006.
- Famiglietti, J. S.: The global groundwater crisis, *Nature Clim Change*, 4, 945–948, <https://doi.org/10.1038/nclimate2425>, 2014.
- Faranda, D., Pascale, S., and Bulut, B.: Persistent anticyclonic conditions and climate change exacerbated the exceptional 2022 European-Mediterranean drought, *Environ. Res. Lett.*, <https://doi.org/10.1088/1748-9326/acbc37>, 2023.
- Füssel, H.-M.: Vulnerability: A generally applicable conceptual framework for climate change research, *Global Environmental Change*, 17, 155–167, <https://doi.org/10.1016/j.gloenvcha.2006.05.002>, 2007.
- Hagenlocher, M., Meza, I., Anderson, C. C., Min, A., Renaud, F. G., Walz, Y., Siebert, S., and Sebesvari, Z.: Drought vulnerability and risk assessments: state of the art, persistent gaps, and research agenda, *Environ. Res. Lett.*, 14, 083002, <https://doi.org/10.1088/1748-9326/ab225d>, 2019.
- IPCC: Climate change 2007: impacts, adaptation and vulnerability: contribution of Working Group II to the fourth assessment report of the Intergovernmental Panel on Climate Change, edited by: Parry, M. L., Cambridge University Press, Cambridge, U.K. ; New York, 976 pp., 2007.
- IPCC: Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press. In Press., 2022.
- IPCC, F., Christopher B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., and IPCC, W. G. I.: Fifth Assessment Report (AR5): Climate Change 2013/2014. Climate change 2014: Impacts, adaptation, and vulnerability ; Part B [Volume 2] [Volume 2, Cambridge University Press, New York, NY, USA, 2014.
- Kim, H., Park, J., Yoo, J., and Kim, T.-W.: Assessment of drought hazard, vulnerability, and risk: A case study for administrative districts in South Korea, *Journal of Hydro-environment Research*, 9, 28–35, <https://doi.org/10.1016/j.jher.2013.07.003>, 2015.



- 537 Medel, K., Kousar, R., and Masood, T.: A collaboration–resilience framework for disaster management supply
 538 networks: a case study of the Philippines, *JHLSCM*, 10, 509–553, [https://doi.org/10.1108/JHLSCM-09-2019-](https://doi.org/10.1108/JHLSCM-09-2019-0066)
 539 0066, 2020.
- 540 Meza, I., Hagenlocher, M., Naumann, G., Vogt, J., and Frischen, J.: Drought vulnerability indicators for global-
 541 scale drought risk assessments: global expert survey results report., Publications Office of the European Union,
 542 Luxembourg, <https://doi.org/10.2760/73844>, 2019.
- 543 Mishra, A. K. and Singh, V. P.: A review of drought concepts, *Journal of Hydrology*, 391, 202–216,
 544 <https://doi.org/10.1016/j.jhydrol.2010.07.012>, 2010.
- 545 Rossi, L., Wens, M., De Moel, H., Cotti, D., Sabino Siemons, A., Toreti, A., Maetens, W., Masante, D., Van
 546 Loon, A., Hagenlocher, M., Rudari, R., Naumann, G., Meroni, M., Avanzi, F., Isabellon, M., and Barbosa, P.:
 547 European drought risk atlas., Publications Office of the European Union, Luxemburg, 2023.
- 548 Sebesvari, Z., Renaud, F. G., Haas, S., Tessler, Z., Hagenlocher, M., Kloos, J., Szabo, S., Tejedor, A., and
 549 Kuenzer, C.: A review of vulnerability indicators for deltaic social–ecological systems, *Sustain Sci*, 11, 575–590,
 550 <https://doi.org/10.1007/s11625-016-0366-4>, 2016.
- 551 Sivakumar, M. V. K., Stefanski, R., Bazza, M., Zelaya, S., Wilhite, D., and Magalhaes, A. R.: High Level meeting
 552 on national drought policy: Summary and major outcomes, *Weather and Climate Extremes*, 3, 126–132,
 553 <https://doi.org/10.1016/j.wace.2014.03.007>, 2014.
- 554 Stenfors, E., Blicharska, M., Grabs, T., and Teutschbein, C.: Droughts in forested ecoregions in cold and
 555 continental climates: A review of vulnerability concepts and factors in socio-hydrological systems, *WIREs Water*,
 556 11, e1692, <https://doi.org/10.1002/wat2.1692>, 2024a.
- 557 Stenfors, E., Blicharska, M., Grabs, T., and Teutschbein, C.: User-Validated Drought Vulnerability Factors in
 558 Forested Cold Climates: Multi-Sectoral Perspectives from Sweden, [https://doi.org/10.5194/egusphere-2024-](https://doi.org/10.5194/egusphere-2024-1988)
 559 1988, 14 August 2024b.
- 560 Teutschbein, C., Quesada Montano, B., Todorović, A., and Grabs, T.: Streamflow droughts in Sweden:
 561 Spatiotemporal patterns emerging from six decades of observations, *Journal of Hydrology: Regional Studies*, 42,
 562 101171, <https://doi.org/10.1016/j.ejrh.2022.101171>, 2022.
- 563 Turner, B. L., Kasperson, R. E., Matson, P. A., McCarthy, J. J., Corell, R. W., Christensen, L., Eckley, N.,
 564 Kasperson, J. X., Luers, A., Martello, M. L., Polsky, C., Pulsipher, A., and Schiller, A.: A framework for
 565 vulnerability analysis in sustainability science, *Proc. Natl. Acad. Sci. U.S.A.*, 100, 8074–8079,
 566 <https://doi.org/10.1073/pnas.1231335100>, 2003.
- 567 UNDRR: Sendai Framework for Disaster Risk Reduction 2015–2030, 2015.
- 568 UNDRR: Global Assessment Report on Disaster Risk Reduction 2019, United Nations, 2019.
- 569 UNDRR: Special report on drought 2021, United Nations Office for Disaster Risk Reduction, Geneva, 173 pp.,
 570 2021.
- 571 Van Loon, A. F.: Hydrological drought explained, *WIREs Water*, 2, 359–392, <https://doi.org/10.1002/wat2.1085>,
 572 2015.
- 573 Wilhite, D. A., Sivakumar, M. V. K., and Pulwarty, R.: Managing drought risk in a changing climate: The role of
 574 national drought policy, *Weather and Climate Extremes*, 3, 4–13, <https://doi.org/10.1016/j.wace.2014.01.002>,
 575 2014.
- 576